



25 November 2010
EMA/HMPC/833398/2009
Patient Health Protection
Committee on Herbal Medicinal Products (HMPC)

Reflection paper on the necessity of initiatives to stimulate the conduct of clinical studies with herbal medicinal products in the paediatric population

Draft

Agreed by Working Party on Community Monographs and Community List (MLWP)	January 2010 March 2010 May 2010
Adoption by HMPC for transmission to PDCO	6 May 2010
Agreed by Working Party on Community Monographs and Community List (MLWP)	November 2010
Adoption by HMPC for release for consultation	25 November 2010
End of consultation (deadline for comments)	<i>15 April 2011</i>
Agreed by Working Party on Community Monographs and Community List (MLWP)	
Coordination with PDCO	
Adoption by HMPC	

Comments should be provided using this [template](#) to hmpc.secretariat@ema.europa.eu

Keywords	HMPC; Herbal medicinal products; clinical studies in paediatric population
----------	--



Reflection paper on the necessity of initiatives to stimulate the conduct of clinical studies with herbal medicinal products in the paediatric population

Table of contents

1. BACKGROUND	3
2. PURPOSE	4
3. DISCUSSION	4
Importance of sound evidence.....	4
State of the art of HMPC monographs.....	5
4. CONCLUSIONS	5
5. REFERENCES	6
6. ANNEX	9
Table 1	9
Table 2	10

1. BACKGROUND

It is well documented that herbal medicinal products (HMPs) are widely used in the general population and specifically in children even if there are important differences among European countries due to specific historical developments and traditions (1-7).

Probably the most important reason for this general popularity is that parents consider them as less dangerous than "conventional" medicinal products because they are "natural", used over hundreds of years and may not be considered as "real drugs".

They can usually be bought without consulting a doctor. Even if consulted, the clinician has documentation on some of the properties of the herbal medicines, but very little clinical information for properly evaluating indications, posology, length of treatment and safety in children.

HMPs are used in children and adolescents for minor but common problems such as Upper Respiratory Tract Infections (URTIs), gastrointestinal disorders, skin problems, sleep disorders, loss of appetite, urinary tract and gynecology disorders (8-13). Moreover there are an increasing number of publications (14-29) regarding the HMPs used, often together with conventional medicinal products, for chronic diseases such as Attention-Deficit-Hyperactivity Disorder (ADHD), depression, inflammatory bowel disease, cystic fibrosis, rheumatoid arthritis, asthma or cancer. This creates the possibility that a HMP may interact with a standard treatment and highlights the need for more information about the use of such therapies.

Directive 2004/24/EC (30) aims to harmonize the market for HMPs and provides a legal basis to facilitate their authorization/registration in Europe. Important tools in the harmonization process are the List of herbal substances, preparations and combinations thereof for use in Traditional Herbal Medicinal Products (THMPs) published by the European Commission and the Community herbal monographs for HMPs having well-established use (WEU) and/or traditional use (TU), established by the Committee on Herbal Medicinal Products (HMPC). Well-established use HMPs have a recognized efficacy and an acceptable level of safety and, usually, have been authorized for more than 10 years in a Member State. THMPs have been in medicinal use for more than 30 years, have been proved to be not harmful in the specified conditions of use and their pharmacological effects or efficacy are plausible on the basis of long-standing use and experience.

Article 8(3)(j) of Directive 2001/83/EC and Article 6(1) of Regulation (EC) 726/2004 (31) require that in order to obtain a marketing authorization, a Summary of Product Characteristics (SmPC) in accordance with Article 11 of Directive 2001/83/EC must be included in the application. The SmPC guideline (32) provides advice on the principles of presenting information in the SmPC. As far as children are concerned, the age limits should reflect the assessment of the available documentation and relate to age intervals where a different dosing is recommended and the information given should relate to ages for which satisfactory efficacy and safety have been shown.

Very often HMPs for children do not completely satisfy the above criteria. This may result in attempts by manufacturers to sell such products as food supplements, so as to overcome the requirements to demonstrate their quality, safety and adequate labelling.

It is unethical that children do not have access to properly assessed medications. For conventional drugs Regulation(EC) No 1901/2006 as amended (33), the 'Paediatric Regulation', revolutionized the regulatory environment for paediatric medicines in Europe by ensuring that medicines for children are of high quality, ethically researched and authorised appropriately, without subjecting children to

unnecessary trials. However, THMPs and HMPs authorised through the well-established medicinal use procedure are not subject to the requirement set out in this legislation to present either studies in the paediatric population in accordance with an agreed Paediatric Investigation Plan or proof of having obtained a waiver or deferral at the time of filing.

It is important to note that, despite such lack of data, a considerable number of European children take HMPs along with or without conventional medicines, so it is important that they are also studied in this age group.

One of the most important aims of the Paediatric Regulation is to reduce the very frequent off-label use of drugs in children, but the situation of HMPs is similar to the off-label use: they are commonly used but have not been adequately studied, they have been on the market for many years via multiple licence-holders, they have no protected intellectual property rights and yet they may be of therapeutic value to children. Moreover performing proper research without any incentives is very costly.

Taking into account the differences between conventional drugs and HMPs it would be useful to improve the situation to ensure medicinal products intended for use in children have been properly assessed in that patient population.

2. PURPOSE

The aim of this document is to highlight the lack of studies on herbal medicine in children and the need for initiatives to stimulate the conduct of clinical studies with HMPs properly designed for children.

3. DISCUSSION

Importance of sound evidence

It is now well accepted that to find the most appropriate treatment for a patient it is necessary to integrate the best evidence available to the clinician with the wishes of the patient. This is important for conventional, complementary and alternative medicines (34-35).

Considering the 'best evidence', the guideline, EMEA/HMPC/104613/2005, on the assessment of clinical safety and efficacy in the preparation of Community herbal monograph (36) refers to the level of evidence and the grading of recommendations used in the WHO General Guidelines for Methodologies on Research and Evaluation of Traditional medicine (37) which considers the strongest evidence is that obtained from meta-analysis of randomized controlled trials, and the weakest that obtained from experts' opinions.

Considering experts' opinions, lack of agreement between them has often been reported (38) raising difficulties for the clinician who has to make a decision on the best treatment for the patient. Regarding the need for information based on evidence, it is difficult to find good quality studies especially in children (39-40) even though many herbal preparations are standardized and can be adequately studied (41).

For this reason, tools to design good trials for HMPs have been proposed by the CONSORT (Consolidated Standards of Reporting Trials) group (42). However, rigorous research is not limited to randomized clinical trials, which also have disadvantages such as costs (both of time and of money) and sometimes ethical problems (43) as well as the risk of incorrect conclusions due to badly designed studies (44).

In some situations observational studies can have advantages (45-46), provided that such studies use validated tools such as the Newcastle-Ottawa Quality Assessment Scale (46). Moreover specific post marketing surveillance studies to define the long-term safety of herbal medicines are the most useful ones (47).

State of the art of HMPC monographs

Seventy-six monographs on HMPs for 155 indications have been published by November 2010.

There are 128 indications for traditional use and 27 for well established use and only one indication (the traditional use for skin disorders and minor wounds of *Avenae fructus*) does not have any age restriction.

The tables in the annex report the results of the analysis.

The SmPC guideline (32) says that a paediatric indication may not be approved if it is not relevant, contraindicated, or because of lack of data or limited/no experience.

Thus, suitable indications for each age group of the tables are considered after excluding those not relevant, contraindicated or generally not recommended and the data show very clearly how the younger the child, the less is the probability of finding indications.

Table 1 show the situation of WEU by age where almost all the indications are approved for adolescents, 56.2% for children older than 6 years and none for the younger ones.

In the case of TU (table 2), 55.9% are those approved for adolescents, around 15% for children from 4 to 12 years, 1.7% from 2 to 4 years and 0.9% for those <2 years of age. These results are mainly due to lack of efficacy and safety data in these age groups.

4. CONCLUSIONS

In spite of frequent use, clinical studies with HMPs in children and adolescents are lacking. For this reason, for the majority of the monographs on HMPs published to date it was not possible to propose any indication for children.

The HMPC considers that there is a need for initiatives to specifically stimulate research in this field to allow the correct use of HMPs in the paediatric population.

The following approaches are proposed:

- 1- Identification of herbal substances/herbal preparations for which a therapeutic benefit is expected (HMPC and PDCO should identify appropriate criteria to select them).
- 2- Provision of guidelines and recommendations for developing appropriate paediatric studies for herbal medicinal products.
- 3- Promotion of funding to collect more data on monitoring safe use in children and to promote further research.

The HMPC would welcome information from stakeholders on experiences with studies on the use of HMPs including THMPs in the paediatric population.

5. REFERENCES

1. Hümer M, Scheller G, Kapellen T, Gebauer C, Schmidt H, Kiess W. Use of herbal medicine in German children - prevalence, indications and motivation *Dtsch Med Wochenschr.* 2010,135(19): pg 959-64.
2. Vlieger AM, van de Putte EM, Hoeksma H. The use of complementary and alternative medicine in children at a general paediatric clinic and parental reasons for use. *Ned Tijdschr Geneesk.* 2006, 150(11): pp 625-30.
3. Madsen H, Andersen S, Nielsen RG, Dolmer BS, Høst A, Damkier A. Use of complementary/alternative medicine among paediatric patients. *Eur J Pediatr.* 2003, 162(5): pp 334-41.
4. Menniti-Ippolito F, Gargiulo L, Bologna E, Forcella E, Raschetti R. Use of unconventional medicine in Italy: a nation-wide survey. *Eur J Clin Pharmacol.* 2002, 58(1): pp 61-4.
5. Crawford NW, Cincotta DR, Lim A, Powel CVE. A cross-sectional survey of complementary and alternative medicine use by children and adolescents attending the University Hospital of Wales. *BMC Complementary and Alternative Medicine* 2006, 6: p 16-9
6. Pommier J, Billot L, Mouchtouris A, Deschamps JP, Romero MI, Zubarew T. French adolescent attitudes towards informal care for physical and emotional or relational problems. *Acta Paediatr.* 2002, 91(4): pp 466-74.
7. Crowe S, Lyons B. Herbal medicine use by children presenting for ambulatory anesthesia and surgery. *Paediatr Anaesth.* 2004, 14(11): pp 916-9.
8. Woolf A.D. Herbal Remedies and Children: Do They Work? Are They Harmful? *Pediatrics* 2003, 112; pp 240-6
9. Linde K, Barrett B, Wölkart K, Bauer R, Melchart D. Echinacea for preventing and treating common cold. *Cochrane Database Sys Rev* 2006: 25: CD000530
10. Hofmann D, Hecker M, Völp A. Efficacy of dry extract of ivy leaves in children with bronchial asthma-a review of randomized controlled trials. *Phytomedicine* 2003, 10(2-3): pp 213-20.
11. Carr RR, Nahata MC. Complementary and alternative medicine for upper- respiratory tract infection in children. *Am J Health Syst Pharm.* 2006, 63(1): pp 33-9.
12. Vohra S, Johnston BC, Laycock KL, Midodzi WK, Dhunnoo I, Harris E, Baydala L. Safety and Tolerability of North American Ginseng Extract in the Treatment of Pediatric Upper Tract Infection: A Phase II Randomized, Controlled Trial of 2 Dosing Schedules. *Pediatrics* 2008, 122 (2): pp 402-410
13. Francis AJ, Dempster RJ. Effect of valerian, *Valeriana edulis*, on sleep difficulties in children with intellectual deficits: randomised trial. *Phytomedicine* 2002, 9(4): pp 273-9.
14. Arnold E, Clark CE, Lasserson TJ, Wu T. Herbal interventions for chronic asthma in adults and children. *Cochrane Database Syst Rev.* 2008, (1):CD005989
15. Tsao J. C. I., Zeltzer LK. Complementary and Alternative Medicine Approaches for Pediatric Pain: A Review of the State-of-the-science. *eCAM* 2005, 2(2): pp 149-59
16. Johnston GA, Bilbao RM, Graham-Brown RA. The use of complementary medicine in children with atopic dermatitis in secondary care in Leicester. *Br J Dermatol.* 2003, 149(3): pp 566-71.

17. Hubner WD, Kirste T. Experience with St John's wort (*Hypericum perforatum*) in children 12 years with symptoms of depression and psychovegetative disturbances. *Phytother Res* 2001, 15: pp 367-70.
18. Martel D, Bussi eres JF, Th eor et Y, Lebel D, Kish S, Moghrabi A, Laurier C. Use of alternative and complementary therapies in children with cancer. *Pediatr Blood Cancer*. 2005, 44(7): pp 660-8.
19. Kelly KM, Jacobson JS, Kennedy DD, Braudt SM, Mallick M, Weiner MA. Use of unconventional therapies by children with cancer at an urban medical center. *J Pediatr Hematol Oncol*. 2000, 22(5): pp 412-6.
20. Shaw A, Thompson EA, Sharp D. Complementary therapy use by patients and parents of children with asthma and the implications for NHS care: a qualitative study. *BMC Health Services Research* 2006, 6:76
21. Gerasimidis K, McGrogan P, Hassank, Edward CA. Dietary modifications, nutritional supplements and alternative medicine in paediatric patients with inflammatory bowel disease. *Aliment Pharmacol Ther* 2008, 27(2): pp 155–65
22. McCann LJ, Newell SJ. Survey of paediatric complementary and alternative medicine use in health and chronic illness. *Arch Dis Child* 2006, 91: pp173–4.
23. Vlieger AM, Blink M, Tromp E, Benninga MA. Use of Complementary and Alternative Medicine by Pediatric Patients With Functional and Organic Gastrointestinal Diseases: Results From a Multicenter Survey. *Pediatrics* 2008, 122 (2): pp 446-51;
24. Soh NL, Walter G. Complementary medicine for psychiatric disorders in children and adolescents. *Curr Opin Psychiatry*. 2008, 21(4): pp 350-5.
25. Lyon MR, Cline JC, Totosy de Zepetnek J, Shan JJ, Pang P, Benishin C. Effect of the herbal extract combination *Panax quinquefolium* and *Ginkgo biloba* on attention-deficit hyperactivity disorder: a pilot study. *Psychiatry Neurosci*. 2001, 26(3): pp 221-8.
26. Wong HH, Smith RG. Patterns of complementary and alternative medical therapy use in children diagnosed with autism spectrum disorders. *Autism Dev Disord*. 2006, 36(7): pp 901-9.
27. Cala S, Crismon ML, Baumgartner J. A survey of herbal use in children with attention-deficit-hyperactivity disorder or depression. *Pharmacotherapy*. 2003 Feb; 23(2):222-30.
28. Weber W, Vander Stoep A, McCarty RL, Weiss NS, Biederman J, McClellan J. *Hypericum perforatum* (St John's wort) for attention-deficit/hyperactivity disorder in children and adolescents: a randomized controlled trial. *JAMA* 2008, 299(22): pp 2633-41
29. Akhondzadeh S, Mohammadi MR, Momeni F. *Passiflora incarnata* in the treatment of attention-deficit hyperactivity disorder in children and adolescents. *Therapy* 2005 2(4) pp 609-14
30. Directive 2001/83/EC as amended by Directives 2003/63/EC, 2004/24/EC and 2004/27/EC
31. Regulation (EC) No 726/2004 of the European Parliament and of the Council of 31 March 2004 laying down Community procedures for the authorisation and supervision of medicinal products for human and veterinary use and establishing a European Medicines Agency.
32. SmPC guideline - http://ec.europa.eu/enterprise/sectors/pharmaceuticals/files/eudralex/vol-2/c/smpc_guideline_rev2_en.pdf

33. Regulation (EC) No 1901/2006 of the European Parliament and of the Council of 12 December 2006 on medicinal products for paediatric use and amending Regulation (EEC) No 1768/92, Directive 2001/20/EC, Directive 2001/83/EC and Regulation (EC) No 726/2004.
34. Angell M, Kassirer JP. Alternative medicine—the risks of untested and unregulated remedies. *N Engl J Med*. 1998, 339: pp 839–841
35. Bondurant S, Sox HC. Mainstream and alternative medicine: converging paths require common standards. *Ann Intern Med* 2005, 142: pp 149–50.
36. Guideline on the assessment of clinical safety and efficacy in the preparation of community herbal monographs for well-established and of community herbal monographs / entries to the community list for traditional herbal medicinal products / substances / preparations EMEA/HMPC/104613/2005
37. WHO General Guidelines for Methodologies on Research and Evaluation of Traditional Medicine, WHO/EDM/TRM/2000.1, WHO Geneva, 2000
38. Ernst E, Pittler MH, Wider B. *The desktop guide to Complementary and alternative Medicine. An evidence-based approach.* Edinburgh: Mosby, 2006, pp: XIII-XV
39. Linde K, Jonas WB, Melchart D, Willich S. The methodological quality of randomized controlled trials of homeopathy, herbal medicines and acupuncture. *Int J Epidemiol* 2001, 30 (3) pp 526-31
40. Moher D, Sampson M, Campbell K, Beckner W, Lepage L, Gaboury I, Berman B. Assessing the quality of reports of randomized trials in pediatric complementary and alternative medicine . *BMC Pediatrics* 2002; 2: 2. Epub 2002 Feb 27.
41. Calapai G, Caputi AP. Herbal Medicines: Can We Do Without Pharmacologist? *eCAM* 2007; 4(S1) pp 41–3
42. Gagnier JJ, Boon H, Rochon P, Moher D, Barnes J, Bombardier C for the CONSORT Group. Recommendations for reporting randomized controlled trials of herbal interventions. *J Clin Epidemiol* 2006; 59(11): pp 1134-1149
43. CEBM, Centre for evidence based medicine, <http://www.cebm.net>.
44. Concato J, Shah N, Horwitz RI. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med*. 2000, 342(25): pp 1887-92.
45. Sackett DL, Wennberg JE. Choosing the best research design for each question. *BMJ* 1997; 315: p 1636
46. Newcastle-Ottawa Scale (NOS)
http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm
47. Ernst E, Cohen MH, Stone J. Ethical problems arising in evidence based complementary and alternative medicine. *J Med Ethics* 2004, 30(2): pp 156–9

6. ANNEX

Table 1

AGE	Well-Established Use							
Years	Not recommended	Not relevant	Contra indicated	Total suitable indications	Medical Advise*	Lack/insufficient data	Limited/no experience	N° and (%) of indications in the monographs
>18				27				27 (100%)
12-18	1	3		22	1			22 (95.6%)
6-12	1	4	6	16	1	6	1	9 (56.2%)
4-6	1	4	6	16	1	6	9	0
2-4	1	4	6	16	1	6	9	0
< 2	1	4	7	15	1	5	9	0

**Salicis cortex (medical advise and only in cases other therapies failed, risk of Reye Syndrome)*

Table 2

AGE	Traditional Use							
Years	Not recommended	Not relevant	Contra indicated	Total suitable indications	Medical Advise +/- lack of data	Lack/ insufficient data	Limited/no experience	N° and (%) of indications in the monographs
>18				128				128
12-18	2	5	3	118	12	37	3	66 (55.9%)
6-12	3	6	5	114	19	65	12	18 (15.8%)
4-6	3	6	5	114	20	67	13	14 (12.3%)
2-4	3	6	5	114	24	67	21	2 (1.7%)
< 2	3	6	13	106	25	67	13	1 (0.9%)